

Klein–Gordon Equation for Quark Pairs in Color Superconductor

B.O.Kerbikov

Institute of Theoretical and Experimental Physics
Moscow, Russia

The wave equation is derived for quark pairs in color superconductor in the regime of low density/strong coupling.

During the last five years color superconductivity became a compelling topic in QCD – see the review papers [1]. Grossly speaking, we have a fair understanding of color superconductivity physics in the high density/weak coupling regime. In the low density/strong coupling region the situation is different. Here, the theory faces the well-known difficulties of the nonperturbative QCD and use is made of the models, like NJL, or instanton gas. By low density we mean the quark densities 3–4 times larger than that in the normal nuclear matter. Model calculations show (see [2] and references in [1]) that at such densities the gap equation acquires a nontrivial solution. This was interpreted [1] as the onset of the Bardeen–Cooper–Schrieffer (BCS) regime, i.e., the formation of the condensate of Cooper pairs made of u - and d -quarks. It is known, however, that nonzero value of the gap is only a signal of the presence of the fermion pairs. Depending on the dynamics of the system, on the fermion density, and on the temperature, such pairs may be either stable, or fluctuating in time, may form a BCS condensate, or the dilute Bose gas, or undergo a Bose condensation. The continuous evolution from the BCS regime to the regime of the Bose–Einstein condensation (BEC) is called the BCS–BEC crossover. Such a transition takes place either by increasing the strength of the interaction, or by decreasing the carrier density. The fact that the BCS wave function may undergo a smooth evolution and describe the tightly bound fermion pairs was first noticed long ago

[3–6]. According to [5], the remark that “there exist a transformation that carries the BCS into BE state” was originally made by F.J.Dyson in 1957 (i.e., the same year when the BCS paper [7] was published). The BCS–BEC crossover for quarks was first discussed in [8]. The general description of the crossover for quarks will be given elsewhere.

In the present paper we shall derive an equation for the quark pairs in the low density/stong coupling limit. This equation is obtained directly from the mean-field gap equation. Our derivation follows the work by P. Nozieres and S. Schmitt-Rink [4], who obtained the Schrödinger equation starting from the BCS solution. Quarks in color superconductor are relativistic particles, and therefore we shall arrive to the Klein–Gordon equation.

Our starting point is the expression for the thermodynamic potential for the 2SC superconductor. In the 2SC phase [1] pairing takes place between u - and d -quarks, while s -quark is out of the game until the density increases so that the chemical potential becomes substantially larger than the mass of the s -quark. Here we consider only the $T = 0$ case. The possible coexistence of the chiral and diquark condensates is neglected on the basis of the Anderson theorem [9]. The expression for the thermodynamic potential reads [2, 9]:

$$\Omega(T = 0, \mu; \Delta) = \frac{\Delta^2}{4g} - \frac{2}{\pi^2} \int dq q^2 \left\{ \sqrt{(E_q - \mu)^2 + \Delta^2} + \sqrt{(E_q + \mu)^2 + \Delta^2} + |\mu - E_q| + E_q \right\}, \quad (1)$$

where $E_q = \sqrt{q^2 + m^2}$.

The four-fermion interaction constant g has a dimension m^{-2} . From (1) one obtains the following gap equation:

$$\Delta = \frac{4g}{\pi^2} \int dq q^2 \left(\frac{\Delta}{\mathcal{E}_q} + \frac{\Delta}{\overline{\mathcal{E}}_q} \right), \quad (2)$$

where $\mathcal{E}_q = \sqrt{(E_q - \mu)^2 + \Delta^2}$, $\overline{\mathcal{E}}_q = \sqrt{(E_q + \mu)^2 + \Delta^2}$.

An important remark concerning the structure of Eqs. (1), (2) is due here. We have tacitly assumed that the four-fermion interaction between quarks is point-like. In general case instead of (2) one should write

$$\Delta_p = \frac{4g}{\pi^2} \int dq q^2 V_{pq} \left(\frac{\Delta_q}{\mathcal{E}_q} + \frac{\Delta_q}{\overline{\mathcal{E}}_q} \right). \quad (3)$$

However, the crude approximation (1), (2) is suffice to obtain the general structure of the Klein–Gordon equation. Next, making use of the standard Bogolubov functional, we introduce the wave functions of the quark–quark and antiquark–antiquark pairs

$$\varphi_p = \frac{\Delta}{\mathcal{E}_p}, \quad \chi_p = \frac{\Delta}{\overline{\mathcal{E}}_p}.$$

Then with a little juggling of (2), (3) we obtain the following set of coupled equations for φ_p and χ_p :

$$(\sqrt{p^2 + m^2} - \mu)\varphi_p = \frac{4g}{\pi^2}(1 - 2n_p) \int dq q^2 (\varphi_q + \chi_q), \quad (4)$$

$$(\sqrt{p^2 + m^2} + \mu)\chi_p = \frac{4g}{\pi^2}(1 - 2\bar{n}_p) \int dq q^2 (\chi_q + \varphi_q), \quad (5)$$

where

$$1 - 2n_p = \frac{E_p - \mu}{\mathcal{E}_p}, \quad 1 - 2\bar{n}_p = \frac{E_p + \mu}{\overline{\mathcal{E}}_p}. \quad (6)$$

These two equations may be recasted into a single Klein–Gordon equation following the standard procedure [10]. Let us define $\psi_p = \varphi_p + \chi_p$, and consider the dilute limit $n_p \ll 1, \bar{n}_p \ll 1$.

Then one gets

$$(p^2 + m^2 - \mu^2)\psi_p = \frac{8g}{\pi^2}\sqrt{p^2 + m^2} \int dq q^2 \psi_q. \quad (7)$$

We remind that non-locality in (7) is of a symbolic character as soon as we use the point-like four-fermion interaction.

From (4), (5), and (7) we see that in the dilute strong coupling limit the chemical potential μ plays the role of the eigenvalue of the Klein–Gordon equation. The point $\mu = m - \mathcal{E}_B$ corresponds to negative non-relativistic chemical potential, typical for the “molecular” limit of the BCS–BEC crossover. The phase diagram in the $(n_p/\bar{n}_p, \mu)$ plane has two symmetric branches corresponding to quarks and antiquarks. Note also that the system described by Eqs. (4), (5) and (7) possesses the “exiton-like” instability.

The author is grateful for discussions and remarks from N.O. Agasian, T.D. Lee, E.V. Shuryak, D.T. Son, M.A. Stephanov, and A.M. Tsvelik. Special thanks to T.Yu. Matveeva for the help in preparing the article. Financial support from BNL and Grant Ssc–1774–2003 is gratefully

acknowledged. We thank INT (Seattle) for its hospitality and the Department of Energy for the support during the Workshop INT-04-1.

It is a pleasure and a honor to submit this paper to Yurii Simonov Festschrift.

References

- [1] K. Rajagopal and F. Wilczek, in *At the Frontier of Particle Physics. Handbook of QCD*, Ed. by M. Shifman (World Sci., Singapore, 2001), Vol.3, p.2061; M. Alford, *Ann. Rev. Nucl. Part. Sci.* **51**, 131 (2001); G. Nardulli, *Riv. Nuovo. Cimento* **25**, 1 (2002).
- [2] J. Berges and K. Rajagopal, *Nucl. Phys. B* **538**, 215 (1999).
- [3] D.M. Eagles, *Phys. Rev.* **186**, 456 (1969); A.J. Leggett, in *Modern Trends in the Theory of Condensed Matter* (Springer-Verlag, Berlin, 1980), p.13.
- [4] P. Nozieres and S. Schmitt-Rink, *J. Low. Temp. Phys.* **59**, 195 (1985).
- [5] P.W. Anderson, *Basic Notions of Condensed Matter Physics* (the Benjamin/Cummings Publ., Menlo Park, California, 1983), p.105.
- [6] P.W. Anderson and W.F. Brinkman, in *Proceedings of the 15th Scottish University Summer School in Physics* (Acad. Press, New York, 1975), p.315.
- [7] J. Bardeen, J. Cooper, and J. Schrieffer, *Phys. Rev.* **108**, 1175 (1975).
- [8] B. Kerbikov, hep-ph/0110197; B. Kerbikov, *Phys. of Atom. Nucl.* **65**, 1918 (2002).
- [9] B. Kerbikov, hep-ph/0106324.
- [10] G. Baym, *Lectures on Quantum Mechanics* (W.A. Benjamin, Inc., New York, 1969), p.499.